



## Bacteria DNA Preparation - Solution Kit

Solution based genomic DNA purification from bacteria

Cat. No.	Amount
PP-206S	100 preparations
PP-206L	500 preparations

**For general laboratory use.**

**Shipping:** shipped at ambient temperature

**Storage Conditions:** store at ambient temperature

**Shelf Life:** 12 months

### Description:

Bacteria DNA Preparation Kit is designed for convenient and fast isolation of genomic DNA from gram-positive and gram-negative bacteria samples. The solution based system minimizes DNA fragmentation that may be problematic in spin-column / filtration based methods. Because phenol or chloroform is not used it is safe and does not produce any harmful waste. Solution based genomic DNA purification kits guarantee minimal DNA fragmentation and yield DNA sized up to 150 kb.

### Expected yield:

Yields of genomic DNA will vary from sample to sample depending on the amount, quality and type of material processed. An amount of approx. 40 µg purified DNA per preparation can be expected.

### Content:

Cell Resuspension Solution

Lysozyme (before use, solve in double distilled water to obtain a final concentration of 100 mg/ml) - store at -20 °C

Cell Lysis Solution

RNase A (before use, solve in double distilled water to obtain a final concentration of 4 mg/ml) - store at -20 °C

Protein Precipitation Solution

Washing Buffer (before use, add 96-99 % Ethanol as indicated on the bottle)

DNA Hydration Solution

### To be provided by you:

Isopropanol (2-propanol) >99 %

96-99 % Ethanol

Microtubes 1.5 ml

Heating Block or Water Bath at 37 °C and 65 °C

### Preparation procedure:

Before start, provide >99 % Isopropanol (2-propanol) (not included in the kit).

**For S pack (100 preps):** Add 250 µl dd-water to the Lysozyme tube, 200 µl dd-water to the RNase A tube and 48 ml 96-99 % Ethanol (not included in the kit) to the Washing Buffer bottle.

**For L pack (500 preps):** Add 250 µl dd-water to each Lysozyme tube, 200 µl dd-water to each RNase A tube and 120 ml 96-99 % Ethanol (not included in the kit) to each Washing Buffer bottle.



## Bacteria DNA Preparation - Solution Kit

Solution based genomic DNA purification from bacteria

Buffer	PP-206S 100 preps	PP-206L 500 preps
Cell Resuspension Solution	32 ml	160 ml
Lysozyme (100 mg/ml)	25 mg	5x 25 mg
Cell Lysis Solution	32 ml	160 ml
RNase A (4 mg/ml)	0.8 mg	5x 0.8 mg
Protein Precipitation Solution	11 ml	55 ml
Washing Buffer	add 48 ml Ethanol (final volume 60 ml)	add 120 ml Ethanol to each bottle (final volume 150 ml each)
DNA Hydration Solution	11 ml	55 ml

- Centrifuge at 15,000 g for 1 min (DNA should be visible as a small white pellet).
- Discard the supernatant and drain tube briefly on clean absorbent paper.
- Add 500 µl Washing Buffer and invert the tube several times to wash the DNA pellet.
- Centrifuge at 15,000 g for 1 min.
- Discard the ethanol carefully.
- Air dry at room temperature for 10-15 min.

### 5 DNA Hydration:

- Add 50-100 µl of DNA Hydration Solution to the dried DNA pellet.
- Hydrate the DNA by incubating at 65 °C for 60 min.
- Store the DNA at 4 °C. For long time storage, store the sample at -20 °C or -80 °C.

### 1a Cell Lysis for Gram-Positive bacteria:

- Transfer 1 ml of cultured cells into a 1.5 ml microtube.
- To harvest the cells centrifuge at 15,000 g for 1 min and discard the supernatant.
- Resuspend the cell pellet in 300 µl of Cell Resuspension Solution.
- Add 2 µl of Lysozyme Solution and mix well by inverting.
- Incubate the tube at 37 °C for 60 min with occasional inverting.
- Centrifuge at 15,000 g for 1 min and discard the supernatant.
- Resuspend the pellet in 300 µl of Cell Lysis Solution.

### 1b Cell Lysis for Gram-Negative Bacteria:

- Transfer 1 ml of cultured cells into a 1.5 ml microtube.
- To harvest the cells centrifuge at 15,000 g for 1 min and discard the supernatant.
- Resuspend the pellet in 300 µl of Cell Lysis Solution.

### 2 RNase Treatment:

- Add 1.5 µl of RNase A Solution and mix by inverting.
- Incubate at 37 °C for 15-30 min and cool on ice for 1 min.

### 3 Protein Precipitation:

- Add 100 µl of Protein Precipitation Solution and vortex vigorously for 20-30 sec.
- Centrifuge at 15,000 g for 5 min.

### 4 DNA Precipitation:

- Transfer the supernatant to a clean 1.5 ml microtube containing 300 µl Isopropanol >99 %.
- Mix the sample by inverting gently for 1 min.